

§ 520.1660

milligrams of oxbendazole per kilogram of body weight; for threadworms, 15 milligrams per kilogram of body weight.

(2) *Indications for use.* For removal and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*); small strongyles (species of the genera *Cylicostephanus*, *Cylicocyclus*, *Cyathostomum*, *Triodontophorus*, *Cylicodontophorus*, and *Gyalocephalus*); large roundworms (*Parascaris equorum*); pinworms (*Oxyuris equi*) including various larval stages; and threadworms (*Strongyloides westeri*).

(3) *Limitations.* Administer by stomach tube in 3 to 4 pints of warm water, or by top dressing or mixing into a portion of the normal grain ration. Prepare individual doses to ensure that each animal receives the correct amount. Horses maintained on premises where reinfection is likely to occur should be re-treated in 6 to 8 weeks. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 78119, Nov. 28, 1980, as amended at 47 FR 39812, Sept. 10, 1982; 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1660 Oxytetracycline.

§ 520.1660a Oxytetracycline and carbomycin in combination.

(a) *Specifications.* (1) Oxytetracycline: The antibiotic substance produced by growth of *Streptomyces rimosus* or the same antibiotic substance produced by any other means.

(2) Carbomycin: The antibiotic substance produced by growth of *Streptomyces halstedii* or the same antibiotic substance produced by any other means.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Special considerations.* The quantities of oxytetracycline in paragraph (e) of this section refer to the activity of oxytetracycline hydrochloride and the quantities of carbomycin listed refer to the activity of an appropriate standard.

(d) *Related tolerances.* See §§ 556.110 and 556.500 of this chapter.

(e) *Conditions of use.* It is used as oxytetracycline hydrochloride plus

21 CFR Ch. I (4-1-01 Edition)

carbomycin base in drinking water of chickens as follows:

(1) *Amount.* 1.0 gram of oxytetracycline and 1.0 gram carbomycin per gallon.

(2) *Indications for use.* As an aid in the prevention and treatment of complicated chronic respiratory disease (air-sac infection) caused by *Mycoplasma gallisepticum* and secondary bacterial organisms associated with chronic respiratory disease such as *E. coli*.

(3) *Limitations.* Administer for not more than 5 days; not for use in chickens producing eggs for human consumption; withdraw 24 hours before slaughter.

§ 520.1660b Oxytetracycline hydrochloride capsules.

(a) *Specifications.* The drug is in capsule form with each capsule containing 125 or 250 milligrams of oxytetracycline hydrochloride. Oxytetracycline is the antibiotic substance produced by growth of *Streptomyces rimosus* or the same antibiotic substance produced by any other means.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs and cats for the treatment of bacterial pneumonia caused by *Brucella bronchiseptica*, tonsillitis caused by *Streptococcus hemolyticus*, bacterial enteritis caused by *Escherichia coli*, urinary tract infections caused by *Escherichia coli*, and wound infections caused by *Staphylococcus aureus*.¹

(2) The drug is administered orally to dogs and cats at a dosage level of 25-50 milligrams per pound of body weight per day in divided doses at 12-hour intervals. The drug can be used for continuation of compatible antibiotic therapy following parenteral oxytetracycline administration where rapidly attained, sustained antibiotic blood levels are required. The duration of treatment required to obtain favorable response will depend to some extent on the severity and degree of involvement and the susceptibility of the infectious

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.